



Northern Goshawk Surveys on BLM Lands within the St. Joe Mountains of Northern Idaho

by

Greg S. Draheim, Mary Terra-Berns, Jay F. Shepherd, Katie M. Kuttner,
Jonathan W. Beals, Charles E. Harris and Wayne Melquist



QL
84.2
.L352
no.
98-19
c. 2

#40825584

ID 88050426

QL
84.2
.L352
NO. 98-19
C.2

Final Report

**NORTHERN GOSHAWK SURVEYS ON
BLM LANDS WITHIN THE
ST. JOE MOUNTAINS OF
NORTHERN IDAHO**

BY

**Greg S. Draheim, Mary Terra-Berns, Jay F. Shepherd, Katie M. Kuttner,
Jonathan W. Beals, Charles E. Harris and Wayne Melquist
Nongame and Endangered Wildlife Program
Bureau of Wildlife**

August 1998

**Idaho Department of Fish and Game
600 S. Walnut, P.O. Box 25
Boise, Idaho 83707
Stephen P. Mealey / Director**

**Challenge Cost-Share Project
#1422-D910-A5-0202**

**Idaho Department of Fish and Game
Bureau of Land Management,
Upper Columbia and Salmon-Clearwater District**

**BLM Library
Denver Federal Center
Bldg. 50, OC-521
P.O. Box 25047
Denver, CO 80225**

TABLE OF CONTENTS

ABSTRACT

A cooperative project between the Department of the Interior, Bureau of Land Management (BLM) and the Idaho Department of Fish and Game was conducted to determine the presence or absence of the Northern Goshawk (*Accipiter gentilis*) on BLM lands in northern Idaho. Responses to taped broadcasts of the adult "alarm" and juvenile "wail" calls were recorded. Two hundred thirty-eight calling stations were surveyed between 6 June 1994 and 1 August 1994. Fourteen responses at 8 different calling stations were elicited. Three hundred forty-three calling stations were surveyed between 8 June 1995 and 14 August 1995. Four responses were recorded at 3 different stations. In 1996, 274 calling stations were surveyed between 6 June and 2 August. One response was recorded during the fledgling period. One hundred sixty-eight calling stations were surveyed between 4 June 1998 and 6 August 1998. One response was recorded during the nestling period. No correlation between sex or age and response rate could be determined from the data collected. Different types of responses as related to sex or age did occur in 1994. Seven goshawk territories were documented, 4 in 1994 and 1 each in 1995 and 1996. Of the 7 territories, 4 were probable breeding territories. A response was detected at the same territory in 1994, 1995, and 1998; however, no active nest sites were found.

LIST OF FIGURES

1. Northern goshawk survey area in BLM lands in the Lemhi, Kootenai, and Shoshone Counties, Idaho. 10

2. Complete coverage of survey area between road 101 and survey area with roads and water (D). 17

TABLE OF CONTENTS

ABSTRACT.....	ii
TABLE OF CONTENTS.....	iii
LIST OF TABLES	iii
LIST OF FIGURES	iii
INTRODUCTION	1
STUDY AREA	1
METHODS	1
RESULTS	3
DISCUSSION	4
LITERATURE CITED	7

LIST OF TABLES

1. Northern goshawk sightings and responses in the St. Joe Mountains, Kootenai, Benewah, and Shoshone Counties, Idaho, 1994-1996, 1998	8
2. Wildlife habitat types at calling stations where goshawk responses were detected in the St. Joe Mountains, Kootenai, Benewah, and Shoshone Counties, Idaho, 1994-1996, 1998	9

LIST OF FIGURES

1. Northern goshawk survey area on BLM lands within Benewah, Kootenai, and Shoshone Counties, Idaho	10
2. Complete coverage of survey areas without roads (A) and survey areas with roads and/or trails (B)	11

TABLE OF CONTENTS

ii ABSTRACT

iii TABLE OF CONTENTS

iv LIST OF TABLES

v LIST OF FIGURES

vi INTRODUCTION

vii STUDY AREA

viii METHODS

ix RESULTS

x DISCUSSION

xi LITERATURE CITED

LIST OF TABLES

1 Northern spotted owl response to the 50 log skidder treatment
in the West-Siskiyou National Monument, Oregon, 1988-1992

2 Wildlife habitat types in cutting areas where spotted owl nests
were located in the West-Siskiyou National Monument, Oregon,
1988-1992

LIST OF FIGURES

1 Northern spotted owl survey area in BLM land near Jewell, Oregon,
and Siskiyou National Monument

2 Changes in crown cover of spruce forest with and without
treatment in the West-Siskiyou National Monument, Oregon, 1988-1992

INTRODUCTION

The Northern Goshawk (*Accipiter gentilis*) is frequently found in forested habitats consisting of intermediate to mature or even over-mature stands with a high degree of canopy closure (Bull and Hohmann 1994, Bright-Smith and Mannan 1994). The understory is generally open and parklike with gentle to moderately steep slopes providing a clear flight path to the nest and unobstructed hunting. These forests are subject to structural changes from timber harvests or other stand altering forces such as disease and fire which diminish suitable goshawk habitat. Therefore, identification of nesting territories and management of these areas is important to maximize the reproductive success of the goshawk (Kennedy and Stahlecker 1993).

The protocol by Woodbridge et al. (1992) identified 3 objectives to consider when planning a survey: (1) determining the presence or absence of goshawks within the project area, (2) confirming of residence in known territories, and (3) annual monitoring of occupancy and reproductive success in known territories. Our objective was to determine the presence or absence of territorial goshawks in the survey areas.

STUDY AREA

Areas surveyed for goshawks were distributed among parcels of land administered by the BLM located in Kootenai, Benewah, and Shoshone counties. Some of the areas surveyed are under consideration for a land exchange timber sale (refer to environmental assessments ID060-96-12 and ID060-95-14). Surveys were conducted in potential habitat in the Rochat Peak, Twin Crag, and Masonia quadrangles (Fig. 1). Areas in the Rochat Peak quadrangle had high road densities with gradual slopes. Stands were mostly mixed conifer composed of grand fir (*Abies grandis*), Douglas-fir (*Pseudotsuga menziesii*), western larch (*Larix occidentalis*), and ponderosa pine (*Pinus ponderosa*). The area within the Twin Crag quadrangle was less accessible and had many steep, rocky slopes. The majority of surveys were conducted in the Latour Peak/Twin Crag area and in the Pine Creek drainage. Stands were composed primarily of mountain hemlock (*Tsuga mertensiana*), subalpine fir (*Abies lasiocarpa*), and Douglas-fir. Survey areas within the Masonia quadrangle had the highest density of roads and slopes, ranging from gradual to steep. The stands were mixtures of ponderosa pine/Douglas-fir, Douglas-fir/grand fir/western red cedar (*Thuja plicata*), and mountain hemlock/sub-alpine fir.

METHODS

We used the survey protocol from the U.S. Forest Service (USFS), Pacific Northwest region (Woodbridge et al. 1992). The basic technique was broadcasting taped goshawk vocalizations along pre-established transects. The objective was to produce an audio and/or visual response from breeding adult goshawks or their offspring.

INTRODUCTION

The Northern Goshawk (*Accipiter gentilis*) is frequently found in forested habitats consisting of intermediate to mature or even over-mature stands with a high degree of canopy closure (Hall and Johnson 1974, Bergsma and Mendenhall 1974). The underlying is generally open and parklike with some secondary forest providing a clear flight path to the nest and unobstructed landing. These birds are subject to structural changes from timber harvests or other stand altering forest management practices and the extent to which suitable habitat. Therefore, identification of nesting territories and management of these areas is important to research on the reproductive success of the goshawk (Kilgus and Johnson 1977).

The purpose of this study was to determine the extent of goshawk nesting territories within the forest area planning a timber harvest (Johnson et al. 1977) identified 3 objectives to consider when (1) determine the extent of nesting territories, and (2) determine the extent of nesting territories. The objective was to determine the extent of nesting territories in the study area.

STUDY AREA

Area surveyed for goshawk nesting territories was divided into 100 ha blocks and administered by the BLM in the Northern, Central, and Southern regions. The area surveyed was under management for timber harvest and timber sale to environmental resources. 1000-2000 ha and 1000-2000 ha blocks were considered in forested habitats in the Kootenai, Lake, Clear, and Adams watersheds. 100 ha blocks in the forest were designated as high quality habitat with suitable habitat. Blocks were made based on the composition of forest in (1) forest type, (2) forest structure, (3) forest management, (4) forest ownership, and (5) forest location. The area within the Lake Clear watershed was less accessible and had many steep, rocky slopes. The majority of areas were located in the Lake Clear watershed and in the Lake Clear watershed. Blocks were composed primarily of conifer forest. It was determined that suitable habitat for goshawk nesting territories was primarily in the Lake Clear watershed and in the Lake Clear watershed. The extent of nesting territories was determined by surveying the area and identifying the extent of nesting territories. The extent of nesting territories was determined by surveying the area and identifying the extent of nesting territories. The extent of nesting territories was determined by surveying the area and identifying the extent of nesting territories.

METHODS

We used the survey protocol from the U.S. Forest Service (U.S. Forest Service 1977) to determine nesting territories. The basic technique was to locate nesting territories. The objective was to determine the extent of nesting territories. The objective was to determine the extent of nesting territories. The objective was to determine the extent of nesting territories.

Suitable nesting habitat was determined prior to the establishment of transects by examining U.S. Geological Survey (USGS) topography maps along with vegetative maps of the survey areas (Keay and Garton 1978). Maps generated by geographic information system (GIS) high-lighting areas with canopy cover > 50% were also utilized. Intermediate to mature stands of forest with > 50% degree of canopy closure were selected for survey transects. Another habitat variable considered was distance to streams, since goshawks appear to be associated with them (Bull and Hohmann 1994, Hargis et al. 1994, Younk and Bechard 1994). In 1995, 1996, and 1998, surveys were also conducted in timber sale areas and in areas considered for a land exchange. Survey transects were established only on BLM-administered lands delineated on GIS maps.

Calling stations were set at 300 m intervals on transects that were located 260 m apart. Transects were denoted by township, range, section, and transect number (e.g. T48N R1W Sec. 35 T4). Calling stations were designated by a superscript on the transect number. Unlike the methods used by Joy et al. (1994), transects were established to accomplish logical coverage of the survey area, including the use of roads and trails whenever possible (Woodbridge et al. 1992) (Fig. 2). In 1995, 1996, and 1998, primarily unroaded areas were surveyed, and topography was considered when placing calling stations. Transects were not straight lines, but stations were still at 300 m intervals. Joy et al. (1994) suggested surveys include 2 or more visits to an area. Transects were surveyed where responses were recorded and also in stands with typical goshawk nesting habitat characteristics (closed canopy, mature trees, open understory, etc.), even though responses had not previously been elicited.

Surveys were conducted during the nestling (June - mid July) and fledgling (mid July - August) periods. Using this time frame reduces disturbance during the courtship and nest construction periods, and maximizes response rates (Kennedy and Stahlecker 1993). The adult goshawk "alarm" call was used from early June to mid-July. The juvenile "wail" call was used from mid-July into August. All surveys were conducted between 0800 and 1730 hours. Prior to conducting the surveys, the observer was trained in northern goshawk identification by sight and sound. Survey techniques were demonstrated to insure familiarity with survey equipment and procedures.

At each calling station, the proper call was played by a Sony Walkman cassette player (model WMA53) through a Radio Shack Power horn (model 32-2037) 6 times. The first broadcast was a 60° turn from the direction of travel. The call lasted 10 seconds followed by a 30-second pause. The next 2 turns were 120° in the same direction followed by 30-second pauses. The procedure was repeated once. Surveys were halted during periods of inclement weather (high winds, heavy rain) and during periods of loud disturbance due to automobile or aircraft traffic.

Data were recorded on standardized Forest Service goshawk survey protocol forms. Sex and age of goshawks were determined when possible. Goshawks were tracked for short distances following visual responses. Nest search forms were completed regardless of search outcome.

RESULTS

1994: Two hundred thirty-eight calling stations were completed between 6 June and 1 August. Fourteen individual goshawk responses were recorded at 9 different calling stations (Table 1). Four of the 9 calling stations had 2 birds responding (Table 1). Responses to the observer ranged from strictly audio with no visual detection, to only audio responses followed by visual detection. An audio response occurred in all 14 goshawk responses. Five goshawk responses were audio/visual while 9 were audio (Table 1).

Six nest searches were conducted following visual detections of adult and juvenile goshawks. No nests were located during these surveys. Goshawk responses were detected at 2 calling stations more than once. Also, 2 goshawks responding at the same time were recorded at 4 calling stations. The age of the responding goshawk was identifiable in all but 2 of 14 detections (Table 1). More detections occurred during the nestling period (12 responses, 153 stations) versus the fledgling period (2 responses, 85 stations) (Table 1).

Douglas-fir was present in 6 of the 8 stands where goshawks were detected (Table 2). Western larch was present in 2 stands, while mountain hemlock and ponderosa pine occurred in 1 stand each (Table 2). One stand was composed of western red cedar and western hemlock (*Tsuga heterophylla*). All stands were intermediate to mature growth with some degree of canopy closure. Although no nests were located, we believed the nesting territories were located in Douglas-fir stands.

1995: Three hundred forty-three calling stations were surveyed between 8 June and 14 August. Four individual goshawk responses were recorded at 3 different calling stations (Table 1). One of the 3 stations had 2 responding goshawks (Table 1). Three of the 4 goshawks were detected by audio/visual responses. The fourth and last goshawk detected, the only observation during the fledgling period, was only detected visually. One hundred ninety-seven calling stations were surveyed during the nestling period and 146 during the fledgling period. Ninety stations surveyed during the nestling period were areas within a land exchange project, 73 were in a timber sale area, and 34 were general surveys in previously determined suitable nesting habitat. All 146 stations surveyed during the fledgling period were in previously determined suitable nesting habitat.

1996: Two hundred seventy-four calling stations were surveyed between 6 June and 2 August. One response was recorded just northeast of the Twin Crags lookout (Table 1). This individual was detected visually and did not vocalize. Detection occurred during the fledgling period.

On 2 separate occasions we observed goshawks while en route to survey locations. The first observation was on 30 July when a goshawk soared from ridge to ridge, crossing Latour Creek road 0.8 km north of the Rochat Peak/Twin Crag road junction. On August 1st, a goshawk was observed on the west side of Latour Peak flying east along the ridge toward Twin Crag. The bird appeared to be hunting and was observed twice. Open areas of grass and rock along the ridge tops support abundant prey species and therefore provide excellent hunting grounds for goshawks and other raptors. It is possible that the goshawk detected at a nearby calling station and the goshawk observed in the area are the same bird. The flying distance between the Latour Creek road and the Latour Peak/Twin Crag area is approximately 4.8 km.

Two hundred six calling stations were surveyed during the nestling period and 68 during the fledgling period. All land exchange and proposed timber sale areas were surveyed during the nestling period (119 calling stations, covering approximately 738 hectares). In the general survey area, approximately 1,822 hectares were surveyed.

1998: One hundred sixty-eight calling stations were surveyed between 4 June and 6 August. Ninety-eight of the calling stations were surveyed during the nestling period and 70 during the fledgling period. The one response recorded during the nestling period (9 June) was on the Hunter/Trapper road off of the East Fork of Pine Creek (Table 1). This was a historical territory that was first found in 1994 and again in 1995 and 1998. The goshawk responded to the "alarm" call twice and then retreated into the forest. A nest search was conducted, but it was not found and the goshawk was never seen. In past surveys both adult birds had been seen (Table 1). There was no response during a subsequent survey of the territory during the fledgling period (15 July). The stand consisted of Douglas-fir/western larch intermediate-sized trees (Table 2) and a somewhat cluttered understory of shrubs and blow-down. Slope was between 20-25%, with an eastern aspect, and a stream was approximately 600 m away.

DISCUSSION

No differences in response rates between adult males and females could be shown from the data collected. In 1994, possible differences existed in the type of response elicited between sexes. During the nestling period males tended to respond from longer distances than females. This may be due in part because males have a much larger home range than females (Woodbridge et al. 1992, Kennedy et al. 1994). Females tended to respond for greater periods of time and in a more aggressive manner (Woodbridge et al. 1992, Joy et al. 1994). This is likely a result of the female not wishing to leave the nest while the unfledged young were still present. One observation of an adult female making a silent fly-by did occur, but only after responding for several minutes from a stationary location.

All juvenile responses were vocal. In 1994, prior to the fledgling period, 2 audio responses by juveniles were recorded (T47N R1E Sec. 12 T3(1) and T48N R1W sec. 28 T4(2); Table 1). Both vocalizations were followed by an adult alarm response. No sightings of juveniles occurred since the birds had not yet fledged and no nests were found. In all cases, the sex could not be determined without a visual detection. The 2 juveniles detected audio/visually were at the same calling station (T48N R2E Sec. 26 T2(1); Table 1) during the fledgling stage. The juveniles continued to vocalize for an extended period of time. The juveniles were reluctant to leave the area even though the observer was in close proximity. The birds were likely recent fledglings and were remaining in the nest territory (Kennedy et al. 1994).

More responses were detected during the nestling stage with the "alarm" call than during the fledgling stage with the "wail" call during the 1994 and 1995 fieldwork. The larger number of stations surveyed during the nestling period in both years may explain this difference. The greater response rate during the nestling stage in 1995 occurred despite land exchange and timber sale areas, which were typically unsuitable nesting habitat. The greatest number of responses, 62%, occurred in stands dominated by Douglas-fir (Table 2). This may be a result of the amount of canopy closure within these stands and their overall maturity. Several stands surveyed in 1996 contained mature trees and had adequate canopy closure however; Douglas-fir was present in small amounts or absent altogether.

The primary tree species in many of the survey stands were mountain hemlock and sub-alpine fir. Morphologically these tree species do not provide an appropriate branching structure for supporting goshawk nests. Although some branches extend horizontally from the bole, most of the branches slope downward, thus precluding large nest structures from being positioned there. Stands of predominantly mountain hemlock and sub-alpine fir may not provide suitable goshawk nesting habitat.

Responses of other raptors were recorded when they occurred. Red-tailed hawks (*Buteo jamaicensis*), Cooper's hawks (*Accipiter cooperii*), sharp-shinned hawks (*Accipiter striatus*), and Barred owls (*Strix varia*) were identified. The incident rate was not high enough to make any correlation between broadcast goshawk vocalizations and the response of other raptors in terms of sex and/or age.

The Woodbridge et al. (1992) method proved to be an effective technique to locate northern goshawks. However, to conduct goshawk surveys and nest searches more effectively, we would recommend that more than 1 surveyor be used. Having only 1 surveyor limited the amount of area that could be surveyed. Although the area to be surveyed for this project was more than 50 percent completed in 1995 (easy areas were completed first, difficult areas were left undone), the remoteness and difficult access of the remaining area to be surveyed suggested the project was roughly 30-40 percent completed. At the conclusion of the 1996 survey season, 70-80 percent of the general survey area was completed. Unsurveyed units were then evaluated using aerial photographs and judged to be highly unlikely to harbor goshawks due to steep slopes with fairly open canopies and lush undergrowth.

All juvenile responses were vocal. In 1994, prior to the fledging period, 2 adults
responses by juveniles were recorded (T47M RJE Sec. 12 T311 and T48N RW Sec. 28 T421;
Table 1). Both vocalizations were followed by an adult alarm response. No change of
juvenile occurred since the birds had not yet fledged and no nest were found. In all cases, the
nest could not be determined within a visual detection. The 2 juveniles detected subsequently
were at the same calling station (T47M RJE Sec. 12 T311; Table 1) during the fledging stage.
The juveniles continued to vocalize for an extended period of time. The juveniles were reluctant
to leave the nest even though the observer was in close proximity. The birds were likely to
fledging and were remaining in the nest during (Kenny et al. 1994).

More responses were observed during the fledging stage with the "alarm" call than during
the fledging stage with the "wail" call during the 1994 and 1995 seasons. The larger number
of responses occurred during the fledging period in both years may explain the difference. The
greater response rate during the fledging stage in 1995 occurred during band exchanges and other
calls were, which were typically associated with nesting behavior. The greater number of responses
observed in 1995 is similar to that reported by Douglas (1994). This may be a result of the amount
of energy spent during these calls and their overall intensity. Several studies surveyed in 1994
conducted near a nest and had a more intense energy than observed. Douglas (1994) was present in
this manner in several situations.

The present study was in many of the ways, which were recorded between the nest and sub-
strate. Although the birds were present in the nest, the birds did not appear to be nesting. Although
the birds were present, they were not nesting. Although the birds were present, they were not nesting.
of the birds, the birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.

Responses of other birds were recorded when they were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.

The # of birds that were recorded when they were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.
The birds were not nesting. The birds were not nesting. The birds were not nesting.

The 1998 field season concluded the survey for goshawks in the St. Joe Mountains of northern Idaho. The area surveyed was approximately 14,245 ha. The only area not searched was a portion of the Latour Creek drainage in the Rochet Peak quadrangle, which amounts to less than 5% of all BLM lands in this area. Five territories were found during the 4 years of the survey. Although no nests were found, 4 of 5 sites had either a pair of goshawks defending the area or juveniles were present indicating active breeding territories. Two calling stations in 1994 (T48N R1W Sec. 28 T4(2) and T47N R1E Sec. 12 T3(1); Table 1) had both adult and juvenile audio responses during the nestling period. An adult audio response was heard from calling station (T48N R1W Sec. 28 T4(2)) on 2 separate occasions. Also in 1994, 2 adult audio vocalizations were detected at calling station T47N R2E Sec. 8 T1(1)(Table 1). A pair of goshawks were detected audio/visually at a nearby calling station in 1995 (T47N R2E Sec. 08 T4⁵; Table 1). Although no nest was found after an extensive search, the highly aggressive behavior of the pair suggests the 2 adults were a breeding pair. This area was surveyed again in 1996 during the fledgling period. Although no response was elicited, an unsuccessful nest search was conducted in the vicinity of last year's detection. In 1998 an audio response was elicited in the same area as 1994 (T47N R2E Sec. 08 T1¹, Table 1). This territory had goshawks on it 2 out of 4 years surveyed (Table 1). In 1994, a visual detection of 2 juveniles at station T48N R2E Sec. 26 T2(2)(Table 1) during the fledgling period indicated that a nest site may have been nearby.

During the 4 years of this survey work, 20 goshawk responses were detected from 1,023 calling stations, resulting in 0.5 responses/100 stations surveyed. This is significantly lower than the 8.0 responses/100 stations that Beals and Harris (1994, 1996) reported while conducting goshawk surveys on the Sawtooth National Forest in southern Idaho. The St. Joe Mountains study area may not have the most suitable forest structure, large trees and a clear understory, that are usual characteristics of goshawk habitat. Most of this area was severely burned in 1910. The remaining large stands of trees were logged for timber or cut for use by the mining industry. Additionally, this area was hit hard by an ice storm in 1996 that caused small to intermediate-sized trees to fall over. These factors combine to create a forest structure generally incompatible with goshawk habitat requirements. There are probably other variables also at work that are beyond the scope of this study.

LITERATURE CITED

- Beals, J. W., and C. E. Harris. 1994. Northern goshawk surveys in the Twin Falls and Burley Ranger Districts of the Sawtooth National Forest. Unpubl. Rep., Idaho Fish and Game, Boise, ID. 17pp.
- _____, and _____. 1996. Northern goshawk surveys in the Twin Falls Ranger District of the Sawtooth National Forest. Unpubl. Rep., Idaho Fish and Game, Boise, ID. 21pp.
- Bull, E. L. and J. H. Hohmann. 1994. Breeding biology of northern goshawks in northeastern Oregon. *Studies in Avian Biology* 16:103-105.
- Bright-Smith, D. J., and R. W. Mannan. 1994. Habitat use by breeding male northern goshawks in northern Arizona. *Studies in Avian Biology* 16:58-65.
- Hargis, C. D., C. McCarthy, and R. D. Perloff. 1994. Home ranges and habitats of northern goshawks in eastern California. *Studies in Avian Biology* 16:66-74.
- Joy, S. M., R. T. Reynolds, and D. G. Leslie. 1994. Northern goshawk broadcast surveys: hawk response variables and survey costs. *Studies in Avian Biology* 16:24-30.
- Keay, J. K., and E. O. Garton. 1978. Terrestrial wildlife habitat inventory of The Rochat study area, BLM, northern Idaho. Final Report. College of Forestry, Wildlife and Range Sciences, University of Idaho. 237pp.
- Kennedy, P. L., and D. W. Stahlecker. 1993. Responsiveness of nesting Northern Goshawks to taped broadcasts of 3 conspecific calls. *J. Wildl. Manage.* 57:249-257.
- Kennedy, P. L., J. M. Ward, G. A. Rinker, and J. A. Gessaman. 1994. Post-fledging areas in northern goshawk home ranges. *Studies in Avian Biology* 16:75-82.
- Scott, J. M., F. Davis, B. Csuti, R. Noss, B. Butterfield, C. Groves, H. Anderson, S. Caicco, F. D'Erchia, T. C. Edwards, Jr., J. Ulliman, and R. G. Wright. 1993. Gap analysis: A geographical approach to protection of biological diversity. *Wildl. Monogr.* 123. 41pp.
- Woodbridge, B., G. Silovsky, K. Austin, C. Puchy, and D. Hays. 1992. Survey protocol for Northern Goshawk on national forest lands in the Pacific Northwest region. 12pp.
- Younk, J. V., and M. J. Bechard. 1994. Breeding ecology of the Northern Goshawk in high-elevation aspen forests of northern Nevada. *Studies in Avian Biology* 16:119-121.

LITERATURE CITED

Baird, J. W., and C. E. Hanks. 1964. Northern goshawk surveys in the Twin Falls and Butte Ranger Districts of the Sawtooth National Forest. Unpubl. Rpt., Idaho Fish and Game, Boise, ID. 17pp.

_____. 1966. Northern goshawk surveys in the Twin Falls Ranger District of the Sawtooth National Forest. Unpubl. Rpt., Idaho Fish and Game, Boise, ID. 21pp.

East, E. J., and J. H. Johnston. 1964. Breeding ecology of northern goshawks in northeastern Oregon. *Studies in Avian Biology* 16:103-107.

Hightshower, D. J., and R. W. Adams. 1964. Nesting and pre-hatching care of northern goshawks in northern Arizona. *Studies in Avian Biology* 16:58-64.

Hagan, C. D., C. McCarthy, and R. D. Pechot. 1966. Home range and habitat of northern goshawk in eastern California. *Studies in Avian Biology* 18:66-74.

Joy, S. M., S. T. Reynolds, and D. G. Leslie. 1964. Northern goshawk population survey: back response variables and survey error. *Studies in Avian Biology* 16:24-30.

Kay, J. E., and E. O. Gahan. 1974. Terrestrial wildlife habitat inventory of the Bitter root, and BLM, northern Idaho. Final Report, College of Forestry, Wildlife and Range Science, University of Idaho. 227pp.

Kennedy, F. J., and G. W. Stiglander. 1961. Reproductive success of northern goshawks in total population of a mountainous area. *J. Wildl. Manage.* 25:249-257.

Kennedy, F. J., J. M. Ward, G. A. Shaw, and J. A. Gorman. 1964. Post-hatching care in northern goshawk home range. *Studies in Avian Biology* 16:75-82.

Scott, J. M., J. E. Kay, B. E. Kay, H. H. Hunsaker, C. G. Geyer, H. Anderson, S. Carson, F. J. Kennedy, J. E. Johnson, B. J. Johnson, and R. G. Wright. 1963. Goshawk survey: A geographical approach to management of mountainous diversity. *Wildl. Manage.* 27:41pp.

Woodbridge, B. G., R. S. K. Smith, C. H. Hays, and D. Hays. 1962. Survey protocol for northern goshawk in central Idaho: birds in the Twin Falls region. 17pp.

Yank, J. V., and M. E. Hunsaker. 1964. Breeding ecology of the northern goshawk in high-altitude habitats in northeastern Oregon. *Studies in Avian Biology* 16:109-121.

Table 1. Northern goshawk sightings and responses in the St. Joe Mountains, Kootenai, Benewah, Shoshone Counties, Idaho, 1994-1996 and 1998.

Date	Transect	Response	Age	Sex
6/8/94	T48N R1W Sec. 35 [T4 (1)] ^a	A/V	A	M
6/15/94	T48N R1W Sec. 34 [T1 (1&2)] ^a	A/V	A	F
6/16/94	T48N R1W Sec. 33 [T3 (2)] ^a	A/V	A	U
6/20/94	T48N R1W Sec. 28 [T4 (2)] ^a	A	U	U
6/21/94	T48N R1W Sec. 28 [T4 (2)] ^a	A	A	U
7/6/94	T48N R1W Sec. 28 [T4 (2)] ^a	A	A	U
7/6/94	T48N R1W Sec. 28 [T4 (2)] ^a	A	J	U
7/6/94	T47N R1E Sec. 12 [T3 (1)]	A	A	U
7/6/94	T47N R1E Sec. 12 [T3 (1)]	A	J	U
7/11/94	T47N R1E Sec. 08 [T1 (3)]	A	U	U
7/14/94	T47N R2E Sec. 08 [T1 (1)]	A	A	M
7/14/94	T47N R2E Sec. 08 [T1 (1)]	A	A	F
7/27/94	T48N R2E Sec. 26 [T2 (1)]	A/V	J	U
7/27/94	T48N R2E Sec. 26 [T2 (1)]	A/V	J	U
7/13/95	T47N R2E Sec. 09 [T3 (5)]	A/V	A	U
7/14/95	T47N R2E Sec. 08 [T4 (5)]	A/V	A	M
7/14/95	T47N R2E Sec. 08 [T4 (5)]	A/V	A	F
8/4/95	T47N R2E Sec. 24 [T16 (4)]	V	A	U
7/15/96	T47N R1E Sec. 08 [T10 (2)]	V	A	F
6/9/98	T47N R2E Sec. 08 [T1 (1)]	A	U	U

^a Transects located on state lands adjacent to BLM lands. Note: brackets denote transects and station number.

Table 2. Wildlife habitat types at calling stations where goshawk responses were detected in the St. Joe Mountains, Kootenai, Benewah, Shoshone Counties, Idaho, 1994-1996 and 1998.

Calling Station	Habitat Type ^a
T48N R1W Sec. 28 [T4 (2)]	Douglas-fir
T48N R1W Sec. 33 [T3 (2)]	western red-cedar/western hemlock
T48N R1W Sec. 34 [T1 (1)]	ponderosa pine/Douglas-fir
T48N R1W Sec. 35 [T4 (1)]	Douglas-fir
T47N R1E Sec. 08 [T1 (3)]	mountain hemlock
T47N R1E Sec. 12 [T3 (1)]	Douglas-fir/western larch
T48N R2E Sec. 26 [T2 (1)]	Douglas-fir
T47N R2E Sec. 08 [T1 (1)]	Douglas-fir/western larch
T47N R2E Sec. 09 [T3 (5)]	western red-cedar/western hemlock
T47N R2E Sec. 08 [T4 (5)]	grand fir
T47N R2E Sec. 24 [T16 (4)]	Douglas-fir
T47N R2E Sec. 08 [T10 (2)]	mountain hemlock
T47N R2E Sec. 08 [T1 (1)]	Douglas-fir/western larch

^a Wildlife habitat types as defined by Scott et al. 1993.






Fig. 1. Northern goshawk survey area on BLM lands located within Benewah, Kootenai and Shoshone Counties, Idaho.



← PROJECT BOUNDARY



PROJECT BOUNDARY

Fig. 2. Complete coverage of survey areas without roads (A), and survey areas with roads and/or trails (B). Adapted from Woodbridge et al. (1992).

Submitted by: Charles E. Harris

Approved by:

IDAHO DEPARTMENT OF FISH AND GAME

Steve Huffaker
Steve Huffaker, Chief
Bureau of Wildlife

Wayne Melquist
Wayne Melquist
State Nongame Wildlife Manager &
Endangered Species Coordinator



Bureau of Land Management

Idaho State Office
1387 S. Vinnell Way
Boise, Idaho 83709

BLM/ID/PT-99/007+1150